


GRAPEVINE FERTILIZATION



ROOT APPLICATION	ROOT APPLICATION	ROOT APPLICATION	FOLIAR USE	FOLIAR USE	FOLIAR USE	FOLIAR USE	FOLIAR USE	FOLIAR USE	ROOT APPLICATION
- Orosoil®	- Mangusta®	- Mangusta®	- Vertyplus®	- Florset®	- Emosprint®	- Vertyplus®	- Aminosprint®	- Kappabrix®	- Fertildung®
- Citoveg®	- Ecoferro®	- Tiger®	- Vigoramin® N8	- Ironleaf Mn®	- Amino.® Calcio	- Aminosprint®	- Calcio e Magnesio		- Ecoferro®
- Aminosprint® N8	- Leopard®	- Teos®	- Ironleaf Mn®	- Vertyplus®	- Amino.® Calcio e Magnesio				- Mangusta®
	- Superguanoxy®	- Aminosprint® N8		- Vigoramin® Boro	- Florset®				
					- Vertyplus®				
TRAPIANTO	VEGETATIVE REST (WINTER)	VEGETATIVE STAGE (SPRING)	LEAF DISTENTION, TILL FLOWER DIFFERENTIATION	FLOWER BUDS	FLOWERING	FLOWER CLOSURE TO RIPENING	GRAPE DEVELOPMENT	GRAPE DEVELOPMENT TO GRAPE RIPENING	POST HARVEST/AUTUMN

The above mentioned doses are indicative and may change according to climatic characteristics of each area (fertility: physical and biological; rainfall and temperature).