


Baseball®

Formulation based on Calcium and Magnesium
with *Ascophyllum nodosum* Algae

CATEGORY
LIQUID
MICROELEMENTS



Features

- It is a liquid product rich in Calcium and Magnesium. Its particular formulation boasts the presence of *Ascophyllum nodosum* Algae to improve the absorption of the two mesoelements and to have a stimulating and anti-stress effect on the plant.
- The product improves the firmness of the skin of the fruit; on the vine, it greatly limits the problem of rachis desiccation and, on apple trees, it helps solving the problem of bitter pitting.
- The formulation does not contain any form of Nitrogen so that it can also be used on apple trees and in the pre-harvest phases on fruit and vegetables.
- Promotes hardening of the peel.
- Makes up for Calcium and Magnesium deficiencies.
- It does not contain Nitrogen.
- It has a biostimulating and anti-stress effect.

Chemical-physical characteristics

pH (5% dilution)

7

Average content in the substance as such

Calcium Oxide

12%

Magnesium Oxide

4%



Via Vialarga 25, 37050
S. Pietro di Morubio (VR) ITALY-UE
Tel. 0039 045 6969004
Fax 0039 045 6969012

fomet@fomet.it
www.fomet.it
f @ in o t



OUR
CERTIFICATIONS

Baseball®

Formulation based on Calcium and Magnesium
with *Ascophyllum nodosum* Algae

CATEGORY
LIQUID
MICROELEMENTS

Suggested dose and method of use for crop nutrition

Arboreal crops	Usage	kg/ha	Notes
Grapevine	Foliar	3 kg/ha for 3 applications	from peppercorn
Rosaceae		5 kg/ha per application	from post-flowering to the end of fruit enlargement
Stone fruit		3 kg/ha per application	
Actinidia		5 kg/ha per application	
Citrus fruits		4 kg/ha per application	
Olive tree		3 kg/ha per application	
Arboreal crops	Usage	kg/ha	Notes
Solanaceae and Cucurbitaceae	Foliar	3 kg/ha per application	from 30 days after transplanting to harvesting
Strawberry		3 kg/ha per application	
Carrot and chicory		3 kg/ha per application	from 30-40 days after
Brassicaceae		3 kg/ha per application	

The above mentioned doses are indicative and may change according to climatic characteristics of each area (fertility; physical and biological; rainfall and temperature). Also, they should be included in the entire fertilization program. Cover the product with in the soil surface is strongly recommended. Product should not be in contact with plant's roots. The product is for professional use only. Keep out of reach of children and animals.

The analytical data written on packaging follow the instruction of the regulation in force. All data included in this publication are indicative. Fomet reserves the right to change them without prior notice.



Properly dispose of packaging.



Keep in a sheltered dry place.